



# Using Distiller's Dried Grains with Solubles in Poultry Diets

**Dr. Jerry Shurson**

**Professor**

**Department of Animal Science**

**University of Minnesota**




## **An Overview of DDGS Production**

## Production of DDGS

- ◆ Tightly linked to the production of fuel ethanol
- ◆ DDGS is a co-product of the dry milling ethanol production process
- ◆ About 40% of ethanol is produced using dry milling
- ◆ The other 60% of ethanol is produced by wet milling
  - co-products include: corn gluten feed, corn gluten meal, and corn germ meal

### Components of Yellow Dent Corn


Starch	61.0 %
Corn Oil	3.8 %
Protein	8.0 %
Fiber	11.2 %
Moisture	16.0 %



Slide courtesy of Ms. Kelly Davis, CVEC

# Production of DDGS

- ◆ Yeasts and enzymes are used to ferment the starch fraction of corn
- ◆ Ethanol and carbon dioxide are produced
- ◆ Distiller's grain and distiller's solubles are the residues remaining after fermentation
- ◆ These fractions are blended and dried to produce distiller's dried grains with solubles (DDGS)



## Dry-Milling Average Yield Per Bushel




- ◆ Ethanol 4.2 liters
- ◆ DDGS 8.2 kg
- ◆ CO<sub>2</sub> 8.2 kg

Slide courtesy of Ms. Kelly Davis, CVEC



## Map of U.S. Ethanol Plants




## DDGS Production

- ◆ 19 new ethanol plants are currently under construction
- ◆ additional capacity is being added to existing plants
- ◆ DDGS will increase from 3.3 million tonnes in 2000 to 5.5 million tonnes in 2005
  - 66% increase in supply of DDGS

## DDGS Production and Use

- ◆ 3.2 to 3.5 million metric tonnes (MT) of DDGS are produced in North America/year
  - ~ 900,000 MT produced in MN-Dakota region
  - ~ 700,000 MT exported to the EU
  - ~ 2.65 million MT fed in U.S. and Canada
    - ~ 2.58 million MT (80%) fed to ruminants
    - ~ 45,000 MT fed in MN turkey industry
    - ~ 27,000 MT used in swine diets

## Markets for DDGS Produced in North America



## Use of DDGS in Swine and Poultry Diets is Increasing

- ◆ DDGS produced by new Midwestern ethanol is higher in nutrient content and digestibility than DDGS from older plants
- ◆ Increased supply of DDGS has made it more economical to replace some of the corn, soybean meal, and dicalcium phosphate

## The Use of DDGS in Poultry Diets



## Historical Use of DDGS in Poultry Diets

- ◆ Fed at low inclusion rates in U.S. poultry industry for many years
  - High B vitamin content (solubles)
  - Source of unidentified growth/reproduction factors?
  - Positive effect on palatability (Alenier and Combs, 1981)
  - Protein source at higher dietary inclusion levels

## Unidentified Growth or Hatchability Factor

- ◆ Growth response (Couch et al., 1957)
  - 5% DDGS in turkey diets
  - 17-32% improvement in gain
- ◆ Feed preference (Alenier & Combs, 1981)
  - 10% DDGS in chicken layer diets
- ◆ Reproduction improvement (Manley, 1978)
  - 3% DDGS in turkey breeder hen diets
  - improvement in egg numbers and hatch (late lay)



## DDGS As Protein Supplement

- ◆ Limiting amino acids (Parsons et al., 1983)
  - Lysine
  - Tryptophan
  - Arginine (perhaps equally limiting with trp)
  
- ◆ Dietary lysine and energy adjustments are needed with inclusion of DDGS
  - growth
  - feed conversion

### Performance of Broiler Chickens (0-42 days) to DDGS in Diets Adjusted and Not Adjusted for Energy (Waldroup et al, 1981)

DDGS Inclusion Level (%)	BW (g)		Gain/Feed	
	Fixed Energy	Variable Energy	Fixed Energy	Variable Energy
0	1288	1206	.513	.493
5	1237	1227	.518	.505
10	1237	1203	.508	.490
15	1220	1165	.513	.444*
20	1246	1164	.498	.467
25	1247	1096*	.500	.446*

\* Different from control

## Lysine Digestibility of DDGS for Poultry

- ◆ Lower than corn due to drying process
- ◆ Lysine bioavailability = 66%
  - Parsons et al. (1983)
- ◆ Lysine digestibility = 65%
  - NRC (1994)

## Lysine Availability (%)

<b>Source</b>	<b>Lysine Bioavail.</b>	<b>Lysine Digest.</b>
Combs & Bossard (1969)	71-93	----
Parsons (1983)	66	82
Heartland (1998)	----	57

## Ingredient Amino Acids (% of Protein)

AA	SBM	Corn	MBM	Canola	DDGS
M+C	3.0	4.6	2.4	4.3	4.2
Lys	6.2	3.0	5.4	5.5	2.8
Iso	4.3	3.2	3.0	3.6	3.6
Arg	7.2	5.0	6.7	6.0	4.4
Trp	1.5	0.9	0.7	1.5	0.8
Thr	4.0	3.5	3.2	4.2	3.8
Val	4.6	4.8	3.8	4.8	4.8

## Amino Acid Digestibility (% of total)

AA	Corn	SBM	MBM	DDGS
	-----%-----			
Met	97.9	94.3	92.3	88.5
Cys	88.2	-----	84.7	78.4
Lys	86.2	91.8	90.3	78.6
Arg	96.1	93.5	94.0	92.5
Tryp	96.8	93.7	95.1	91.8
Thr	81.1	84.3	90.1	82.5
Iso	86.4	90.9	92.2	89.1
Val	93.3	89.4	90.2	88.1

## Energy Utilization of DDGS by Poultry

- ◆ Crude protein = 27.4%
- ◆ Crude fat = 9%
- ◆ ME, kcal/kg = 2,480
- ◆ ME/GE, % = 49.1
- ◆ ME (% of corn) = 74.1

## Maximum Inclusion Rates of DDGS in Poultry Diets

- ◆ Turkey, pullet, and broiler starter = 5%
- ◆ Turkey and broiler finisher = 15%
- ◆ Pullet developer = 10%
- ◆ Layer (peak) = 10%
- ◆ Layer (late lay) = 10%


Source: ME Jackson, 2002 Midwest Poultry Federation Convention, St. Paul, MN

## Use of DDGS in Poultry Diets

- ◆ High inclusion rates have also provided good results
  - Favorable results with 25% DDGS in broiler diets
    - Waldroup et al., 1981
  - 15% DDGS in layer diets reduces fatty liver incidence
    - Jensen et al., 1974; Jensen, 1987; Akiba et al., 1983
  - 12% DDGS turkey diets gave similar performance to corn-soybean meal diets
    - Noll, 2002

## DDGS Quality is Highly Variable

- ◆ Nutritionists want PREDICTABILITY AND CONSISTENCY in feed ingredients.
- ◆ The keys for getting maximum value from DDGS are:
  - “Know what you have (or want)”
  - and
  - “Know how to use it”



## DDGS Quality is Variable


- ◆ Color ranges from very light to very dark
- ◆ Odor ranges from sweet to smoky or burnt
- ◆ Range in concentration in selected nutrients:
  - Dry matter – 87 to 93%
  - Crude protein – 23 to 29%
  - Crude fat – 3 to 12%
  - Ash – 3 to 6%
  - Lysine – 0.59 to 0.89%

Source: Cromwell et al. (1993)


## Nutrient Profile of Corn Distiller's Dried Grains with Solubles (DM Basis)

Nutrient	MW DDGS	Low Quality DDGS	NRC (1998)
Dry matter, %	88.9	88.3	93.0
Crude protein, %	30.2	28.1	29.8
Fat, %	10.9	8.2	9.0
Fiber, %	8.8	7.1	4.8
Calcium, %	0.06	0.44	0.22
Phosphorus, %	0.89	0.90	0.83
P availability, %	90.0	?	79.0
DE, kcal/kg	3965	3874	3449
ME, kcal/kg	3592	3521	3038
Lys, %	0.83	0.53	0.67
App. Dig. Lys, %	0.44	0.00	0.34
Met, %	0.55	0.50	0.54
App. Dig. Met, %	0.32	0.24	0.42
Thr, %	1.13	0.98	1.01
App. Dig. Met, %	0.62	0.36	0.60
Trp, %	0.24	0.19	0.27
App. Dig Trp, %	0.15	0.15	0.15

## Growth of Chicks Fed Nine Sources of DDGS



## Feed Conversion of Chicks Fed Nine Sources of DDGS



## Nutritional Value of DDGS for Poultry

- ◆ Must use high quality DDGS
  - Golden color = high amino acid digestibility
- ◆ Excellent energy and available phosphorus source
- ◆ Nutritional value higher than previously thought
- ◆ Unidentified growth factors?
  - 5% DDGS resulted in 17-32% improvement in gain
  - 3% DDGS in turkey breeder hen diets increased egg numbers and hatch
- ◆ Effective partial replacement for corn and soybean meal



## Quality Considerations for Selecting DDGS Sources for Swine and Poultry

### ◆ Nutrient Specifications

- Moisture – maximum 12%
- Protein – minimum 26.5%
- Fat – minimum 10%
- Fiber – maximum 7.5%

## Quality Considerations for Selecting DDGS Sources for Swine and Poultry

### ◆ Physical characteristics

- Bulk density – .44 to .48 kg/cubic meter
- Particle size:
  - maximum coarse particles - 10% on 2000 screen
  - maximum fine particles - 15% on 600 screen & in pan
- Smell – fresh, fermented
- Color – goldenrod

## **Maximizing the Value of Corn DDGS in Poultry Diets**

- ◆ Formulate diets using digestible amino acid values
- ◆ High available P reduces the level of dietary P supplementation
- ◆ Adding 5 % DDGS to may improve feed preference, egg number, and hatchability in breeder hens

## **Limitations of Using DDGS in Poultry Diets**

- ◆ Must be golden color and highly digestible
- ◆ High fiber limits its maximum inclusion rate in poultry diets
- ◆ Excess nitrogen can be minimized by using synthetic amino acids

## U of M DDGS Web Site

We have developed a DDGS web site featuring:

- \* research summaries (swine, poultry, dairy, & beef)
- \* presentations given
- \* links to other DDGS related web sites

Visit this web site at:

[www.ddgs.umn.edu](http://www.ddgs.umn.edu)